

Total Teleportation of an Entangled State

Hai-Woong Lee

Department of Physics, Korea Advanced Institute of Science and Technology, Taejon 305-701, Korea

(Dated: February 1, 2008)

We present a setup for a direct, total teleportation of a single-particle entangled state. Our scheme consists of a parametric down-conversion source, which emits a pair of entangled photons, and a dual Bell measurement system with two beam splitters and two pairs of detectors. An entangled state shared by a pair of beams can be directly transferred to another pair of beams with a 50% probability of success. By a straightforward generalization of the scheme, total teleportation of a multi-particle entangled state can also be performed.

PACS number(s): 03.67.-a, 03.65.Ta, 42.50.-p

The recent progress and interest in quantum information science stems largely from the realization that nonclassical features of quantum systems can be utilized to achieve performance in communications and computations that is superior to that based on classical means[1]. The nonlocal nature of quantum systems arising from entanglement, in particular, has played a central role in many recent exciting developments in this area of research. Entanglement is an essential resource for many ingenious applications in quantum information science such as quantum teleportation[2, 3] and quantum cryptography[4, 5] as well as for fundamental studies in quantum mechanics related to Einstein-Podolsky-Rosen (EPR) paradox[6] and Bell's theorem[7]. In Bell's test experiments and in quantum teleportation and cryptography experiments, an important issue is whether one can generate and manipulate entanglement in a controlled way.

In this paper we address ourselves to the issue of entanglement transfer, more specifically, quantum teleportation of an unknown entangled state. As pointed out by Bennett et al.[2] in their original proposal for quantum teleportation, entanglement can be transferred through teleportation of the state of one of the entangled pair. This method, known as entanglement swapping[8, 9, 10], provides partial teleportation of entanglement in the sense that the teleported state is only a part of the entangled state. We propose an alternative method in which the entire entangled state is transferred directly from one place to another. The method is demonstrated below for the case when the state to be teleported is a single-particle entangled state.

The teleportation scheme we propose is depicted in Fig. 1. The state we wish to teleport is a single-photon entangled state of the form $\alpha|1\rangle_{a_1}|0\rangle_{a_2} + \beta|0\rangle_{a_1}|1\rangle_{a_2}$, where α and β are unknown except that $|\alpha|^2 + |\beta|^2 = 1$. Here $|1\rangle$ and $|0\rangle$ refer to a one-photon state and a vacuum state, respectively, and the subscripts a_1 and a_2 represent the modes of photon in the beams a_1 and a_2 , respectively. This state may, for example, be generated when a photon is incident on a beam splitter of unknown reflection and transmission coefficients, $r = \alpha$, and $t = \beta$, as shown in the lower left part of Fig. 1. At the source station we have a parametric down-conversion source which emits a pair of entangled photons. These photons emerge either through the ports A_1 and B_1 or through A_2 and B_2 . Such a source has been considered in past studies of entanglement swapping[8, 9, 10]. The photons are thus in a two-photon entangled state

$$|\psi\rangle_{source} = \frac{1}{\sqrt{2}}(|1\rangle_{A_1}|0\rangle_{A_2}|1\rangle_{B_1}|0\rangle_{B_2} + |0\rangle_{A_1}|1\rangle_{A_2}|0\rangle_{B_1}|1\rangle_{B_2}) \quad (1)$$

The beams A_1 and A_2 are directed to Alice and the beams B_1 and B_2 to Bob.

The state of the entire photon field before being detected by the four detectors of Fig.1 is given by

$$|\psi\rangle_{in} = |\psi\rangle_{source}(\alpha|1\rangle_{a_1}|0\rangle_{a_2} + \beta|0\rangle_{a_1}|1\rangle_{a_2}) \quad (2)$$

At Alice's station the beam A_1 from the source is combined via a 50/50 beam splitter with the beam a_1 for a Bell measurement. At the same time the beam A_2 is combined via another 50/50 beam splitter with the beam a_2 for another Bell measurement. Expressing the state $|\psi\rangle_{in}$ in the Bell basis, we can write

$$\begin{aligned} |\psi\rangle_{in} = \frac{1}{2\sqrt{2}} & [(|\Phi_{A_1a_1}^{(+)}\rangle|\Phi_{A_2a_2}^{(+)}\rangle - |\Phi_{A_1a_1}^{(-)}\rangle|\Phi_{A_2a_2}^{(-)}\rangle)(\alpha|1\rangle_{B_1}|0\rangle_{B_2} + \beta|0\rangle_{B_1}|1\rangle_{B_2}) \\ & + (|\Phi_{A_1a_1}^{(-)}\rangle|\Phi_{A_2a_2}^{(+)}\rangle - |\Phi_{A_1a_1}^{(+)}\rangle|\Phi_{A_2a_2}^{(-)}\rangle)(\alpha|1\rangle_{B_1}|0\rangle_{B_2} - \beta|0\rangle_{B_1}|1\rangle_{B_2}) \\ & + (|\Psi_{A_1a_1}^{(+)}\rangle|\Psi_{A_2a_2}^{(+)}\rangle - |\Psi_{A_1a_1}^{(-)}\rangle|\Psi_{A_2a_2}^{(-)}\rangle)(\beta|1\rangle_{B_1}|0\rangle_{B_2} + \alpha|0\rangle_{B_1}|1\rangle_{B_2}) \\ & + (|\Psi_{A_1a_1}^{(-)}\rangle|\Psi_{A_2a_2}^{(+)}\rangle - |\Psi_{A_1a_1}^{(+)}\rangle|\Psi_{A_2a_2}^{(-)}\rangle)(\beta|1\rangle_{B_1}|0\rangle_{B_2} - \alpha|0\rangle_{B_1}|1\rangle_{B_2})] \end{aligned} \quad (3)$$

where the Bell states are defined by

$$\begin{aligned} |\Phi_{A_i a_i}^{(\pm)}\rangle &= \frac{1}{\sqrt{2}}(|1\rangle_{A_i}|1\rangle_{a_i} \pm |0\rangle_{A_i}|0\rangle_{a_i}) \\ |\Psi_{A_i a_i}^{(\pm)}\rangle &= \frac{1}{\sqrt{2}}(|1\rangle_{A_i}|0\rangle_{a_i} \pm |0\rangle_{A_i}|1\rangle_{a_i}) \end{aligned} \quad (4)$$

If Alice's Bell measurement results in a simultaneous detection of one photon each at the detector D_{11} and D_{21} and no photon at D_{12} and D_{22} (corresponding to the input state $|\Psi_{A_1 a_1}^{(+)}\rangle|\Psi_{A_2 a_2}^{(+)}\rangle$), or a simultaneous detection of one photon each at D_{12} and D_{22} and no photon at D_{11} and D_{21} (corresponding to the input state $|\Psi_{A_1 a_1}^{(-)}\rangle|\Psi_{A_2 a_2}^{(-)}\rangle$), then the state of the photon at Bob's station reduces to $\beta|1\rangle_{B_1}|0\rangle_{B_2} + \alpha|0\rangle_{B_1}|1\rangle_{B_2}$ [11]. If Bob is informed of such a measurement result from Alice, he does not need to perform any unitary transformation. He only needs to identify the beam B_2 as a teleported version of the beam a_1 and the beam B_1 as a teleported version of the beam a_2 , and the teleportation is achieved. If Alice's Bell measurement results in a simultaneous detection of one photon each at D_{11} and D_{22} and no photon at D_{12} and D_{21} (corresponding to the input state $|\Psi_{A_1 a_1}^{(+)}\rangle|\Psi_{A_2 a_2}^{(-)}\rangle$), or a simultaneous detection of one photon each at D_{12} and D_{21} and no photon at D_{11} and D_{22} (corresponding to the input state $|\Psi_{A_1 a_1}^{(-)}\rangle|\Psi_{A_2 a_2}^{(+)}\rangle$), then the state of the photon at Bob's station is $\beta|1\rangle_{B_1}|0\rangle_{B_2} - \alpha|0\rangle_{B_1}|1\rangle_{B_2}$. An appropriate unitary transformation can then be performed by Bob, and the teleportation is achieved. Alice's Bell measurement, however, cannot distinguish between the states $|\Phi_{A_1 a_1}^{(+)}\rangle|\Phi_{A_2 a_2}^{(+)}\rangle$, $|\Phi_{A_1 a_1}^{(-)}\rangle|\Phi_{A_2 a_2}^{(-)}\rangle$, $|\Phi_{A_1 a_1}^{(+)}\rangle|\Phi_{A_2 a_2}^{(-)}\rangle$, and $|\Phi_{A_1 a_1}^{(-)}\rangle|\Phi_{A_2 a_2}^{(+)}\rangle$, and therefore the probability of success for the present scheme of entanglement teleportation is 50%, the same as that for the standard teleportation scheme [2].

The teleportation experiment described above can be performed in an "event-ready" fashion by using the technique of entanglement swapping, as suggested by Zukowski et al. [8]. For this purpose we employ at the source station the setup shown in Fig. 2. It consists of two independent sources, two 50/50 beam splitters and two pairs of detectors. The two sources each emit one pair of entangled photons in the state $\frac{1}{\sqrt{2}}(|1\rangle_{A_1}|0\rangle_{B_2}|1\rangle_{G_1}|0\rangle_{H_2} + |0\rangle_{A_1}|1\rangle_{B_2}|0\rangle_{G_1}|1\rangle_{H_2}) + \frac{1}{\sqrt{2}}(|1\rangle_{A_2}|0\rangle_{B_1}|1\rangle_{G_2}|0\rangle_{H_1} + |0\rangle_{A_2}|1\rangle_{B_1}|0\rangle_{G_2}|1\rangle_{H_1})$. Detection of one photon each at D_{G_1} and D_{H_1} and no photon at D_{G_2} and D_{H_2} , or detection of one photon each at D_{G_2} and D_{H_2} and no photon at D_{G_1} and D_{H_1} , would indicate that the state collapses into $|\Psi_{G_1 G_2}^{(\pm)}\rangle|\Psi_{H_1 H_2}^{(\pm)}\rangle \frac{1}{\sqrt{2}}(|1\rangle_{A_1}|0\rangle_{A_2}|1\rangle_{B_1}|0\rangle_{B_2} + |0\rangle_{A_1}|1\rangle_{A_2}|0\rangle_{B_1}|1\rangle_{B_2})$. Thus, the state of the photons emerging eventually from the source subjected to either of the above detection results becomes exactly the state $|\psi\rangle_{source}$ of Eq. (1). The detectors D_{11} , D_{12} , D_{21} , and D_{22} can be activated when the detectors D_{G_1} , D_{G_2} , D_{H_1} , and D_{H_2} register either of the above results.

The proposed scheme of teleporting single-particle entanglement can be generalized to the case of multi-particle entanglement in a straightforward way. Let us consider teleportation of an N-particle entangled state of the form

$$\alpha|1\rangle_{a_1}|0\rangle_{a_2}|1\rangle_{b_1}|0\rangle_{b_2} \cdots |1\rangle_{p_1}|0\rangle_{p_2} + \beta|0\rangle_{a_1}|1\rangle_{a_2}|0\rangle_{b_1}|1\rangle_{b_2} \cdots |0\rangle_{p_1}|1\rangle_{p_2} \quad (5)$$

The quantum channel that links Alice and Bob is provided by an (N+1)-particle entangled state

$$\frac{1}{\sqrt{2}}(|1\rangle_{A_1}|0\rangle_{A_2}|1\rangle_{B_1}|0\rangle_{B_2} \cdots |1\rangle_{P_1}|0\rangle_{P_2}|1\rangle_{Q_2}|0\rangle_{Q_2} + |0\rangle_{A_1}|1\rangle_{A_2}|0\rangle_{B_1}|1\rangle_{B_2} \cdots |0\rangle_{P_1}|1\rangle_{P_2}|0\rangle_{Q_1}|1\rangle_{Q_2}) \quad (6)$$

In Eqs. (5) and (6) the letters p and P serve to identify the N-th particle and the letter Q the (N+1)-th particle. The (N+1)-particle entangled state of Eq. (6) is generated at the source station. The beams A_1 and A_2 , along with the beams containing the unknown entangled state, are sent to Alice, while all other beams are sent to Bob. With (N+1) beam splitters and (N+1) pairs of detectors, Alice then makes a Bell measurement on the pairs (A_1, a_1) , (A_2, a_2) , (b_1, b_2) , (c_1, c_2) , $\dots$, (p_1, p_2) , separately.

The photon field state given by the product of Eqs. (5) and (6) can be expressed in the Bell basis. Straightforward algebra shows that the photon field state expressed in the Bell basis can be divided into four groups, each associated with a different field state at Bob's station. Two of the four groups contain products of the states of the form $|\Phi_{A_1 a_1}^{(\pm)}\rangle|\Phi_{A_2 a_2}^{(\pm)}\rangle$, and are not of much use, because these states cannot be distinguished from each other by Alice's Bell measurement. The remaining two groups can be expressed as

$$\begin{aligned} &\sum_{even} (\pm) |\Psi_{A_1 a_1}^{(\pm)}\rangle |\Psi_{A_2 a_2}^{(\pm)}\rangle |\Psi_{b_1 b_2}^{(\pm)}\rangle |\Psi_{c_1 c_2}^{(\pm)}\rangle \cdots |\Psi_{p_1 p_2}^{(\pm)}\rangle (\beta|1\rangle_{B_1}|0\rangle_{B_2}|1\rangle_{C_1}|0\rangle_{C_2} \cdots |1\rangle_{Q_1}|0\rangle_{Q_2} \\ &\quad + \alpha|0\rangle_{B_1}|1\rangle_{B_2}|0\rangle_{C_1}|1\rangle_{C_2} \cdots |0\rangle_{Q_1}|1\rangle_{Q_2}) \\ &+ \sum_{odd} (\pm) |\Psi_{A_1 a_1}^{(\pm)}\rangle |\Psi_{A_2 a_2}^{(\pm)}\rangle |\Psi_{b_1 b_2}^{(\pm)}\rangle |\Psi_{c_1 c_2}^{(\pm)}\rangle \cdots |\Psi_{p_1 p_2}^{(\pm)}\rangle (\beta|1\rangle_{B_1}|0\rangle_{B_2}|1\rangle_{C_1}|0\rangle_{C_2} \cdots |1\rangle_{Q_1}|0\rangle_{Q_2} \\ &\quad - \alpha|0\rangle_{B_1}|1\rangle_{B_2}|0\rangle_{C_1}|1\rangle_{C_2} \cdots |0\rangle_{Q_1}|1\rangle_{Q_2}) \end{aligned} \quad (7)$$

where $\sum_{\text{even}} [\sum_{\text{odd}}]$ denotes that, out of all possible combinations of $|\Psi^{(+)}\rangle$'s and $|\Psi^{(-)}\rangle$'s in the product of $(N+1)$ Bell states, only those for which the total number of $|\Psi^{(-)}\rangle$'s is even [odd] should be included in the summation, and $(\pm)$ immediately following the summation sign indicates that some of the terms in the summation are added and some others subtracted.

It is important to note that different combinations of the product of $(N+1)$ $|\Psi^{(\pm)}\rangle$'s have different combinations of detectors registering a photon. Thus, when the $(N+1)$ pairs of detectors register a particular combination of detection coincidence, we know which combination of the product of $|\Psi^{(\pm)}\rangle$'s it corresponds to and consequently which state Bob has in the beams $(B_1, B_2, C_1, C_2, \dots, Q_1, Q_2)$. An appropriate unitary transformation performed by Bob then completes the teleportation process. Since the state given by Eq.(7) represents one half of the entire initial photon state, the probability of success for teleportation of an N -particle entangled state is 50%.

Let us consider the case $N=2$. This case corresponds to teleporting a two-particle entangled state using three-particle entanglement and has recently been considered by Shi et al.[12] and by Marinatto and Weber[13]. Our scheme for teleporting a two-particle entangled state $\alpha|1\rangle_{a_1}|0\rangle_{a_2}|1\rangle_{b_1}|0\rangle_{b_2} + \beta|0\rangle_{a_1}|1\rangle_{a_2}|0\rangle_{b_1}|1\rangle_{b_2}$ is shown in Fig.3. Here the quantum channel should be provided by a three-particle Greenberger-Horne-Zeilinger (GHZ) state[14] $\frac{1}{\sqrt{2}}(|1\rangle_{A_1}|0\rangle_{A_2}|1\rangle_{B_1}|0\rangle_{B_2}|1\rangle_{C_1}|0\rangle_{C_2} + |0\rangle_{A_1}|1\rangle_{A_2}|0\rangle_{B_1}|1\rangle_{B_2}|0\rangle_{C_1}|1\rangle_{C_2})$. This type of GHZ state can be generated using the method of Zeilinger et al.[15] In this situation, a coincidence detection of a single photon by the detectors $(D_1, D_3, \text{ and } D_5)$, or $(D_1, D_4, \text{ and } D_6)$, or $(D_2, D_3, \text{ and } D_6)$, or $(D_2, D_4, \text{ and } D_5)$, collapses the state at Bob's station into $\beta|1\rangle_{B_1}|0\rangle_{B_2}|1\rangle_{C_1}|0\rangle_{C_2} + \alpha|0\rangle_{B_1}|1\rangle_{B_2}|0\rangle_{C_1}|1\rangle_{C_2}$. This occurs with a 25% probability. On the other hand, a coincidence detection of a single photon by the detectors $(D_1, D_3, \text{ and } D_6)$, or $(D_1, D_4, \text{ and } D_5)$, or $(D_2, D_3, \text{ and } D_5)$, or $(D_2, D_4, \text{ and } D_6)$, collapses the state at Bob's station into $\beta|1\rangle_{B_1}|0\rangle_{B_2}|1\rangle_{C_1}|0\rangle_{C_2} - \alpha|0\rangle_{B_1}|1\rangle_{B_2}|0\rangle_{C_1}|1\rangle_{C_2}$. This occurs also with a 25% probability. The case $N=3$ corresponds to teleportation of an unknown GHZ-type entangled state $\alpha|1\rangle_{a_1}|0\rangle_{a_2}|1\rangle_{b_1}|0\rangle_{b_2}|1\rangle_{c_1}|0\rangle_{c_2} + \beta|0\rangle_{a_1}|1\rangle_{a_2}|0\rangle_{b_1}|1\rangle_{b_2}|0\rangle_{c_1}|1\rangle_{c_2}$. The quantum channel in this case is provided by a four-particle entangled state $\frac{1}{\sqrt{2}}(|1\rangle_{A_1}|0\rangle_{A_2}|1\rangle_{B_1}|0\rangle_{B_2}|1\rangle_{C_1}|0\rangle_{C_2}|1\rangle_{E_1}|0\rangle_{E_2} + |0\rangle_{A_1}|1\rangle_{A_2}|0\rangle_{B_1}|1\rangle_{B_2}|0\rangle_{C_1}|1\rangle_{C_2}|0\rangle_{E_1}|1\rangle_{E_2})$, which can be generated, for example, using the method of Bose et al.[9] which is a generalized version of the method of Zukowski et al.[8] or of Zeilinger et al.[15].

Bennett et al.[2], in their original proposal for quantum teleportation, have already noted that teleportation can also be achieved for entangled states. Let particles A and B be an entangled pair produced at the source station and sent to Alice and Bob, respectively, and let particles a and b be a pair in an unknown entangled state. If Alice makes a Bell measurement on the particles A and a , the state of particle a can be teleported to particle B at Bob's station. In addition, Bennett et al. showed that the particles B and b are entangled exactly the same way the particles a and b were before teleportation. This is entanglement swapping[8, 9, 10] and provides a way of teleporting an entangled state via teleportation of the state of only one particle of the entangled pair. Teleportation of an unknown single-particle entangled state using this method has been discussed in our earlier publication[16]. We note, however, that, if Bob is to have the desired entangled pair in his possession, not only the particle B should be sent to him from the source station, but also the particle b should be arranged to be sent to him. In contrast, the scheme described here offers a way for a direct, total teleportation of an entangled state. The entangled state shared by the two beams a_1 and a_2 of Fig. 1 is directly teleported to the beams B_1 and B_2 , which were sent to Bob from the source station. The beams a_1 and a_2 in an unknown entangled state are both subjected to Alice's Bell measurement and neither needs to be sent to Bob. We also note that the present scheme allows teleportation of an entangled state with only three photons, the same number necessary to teleport a qubit of states in the standard teleportation scheme.

One may argue that the necessity of sending the particle b to Bob in the standard teleportation scheme can be avoided if the particle b is subjected to a second Bell measurement at Alice's station. It then is necessary that the source station generates two entangled pairs, say (A_1, B_1) and (A_2, B_2) . The particles A_1 and A_2 are sent to Alice and the beams B_1 and B_2 to Bob. Alice makes two successive Bell measurements, first on the pair $(A_1 \text{ and } a)$ and second on the pair $(A_2 \text{ and } b)$. the entanglement can be transferred this way from the particles a and b to the particles B_1 and B_2 . This method, however, has only a 25% probability of success, because each Bell measurement is associated with a 50% probability of success. Furthermore, the method may not succeed if the quantum channel is noisy, because the initial state for the second Bell measurement is no longer pure[17]. Apparently, the present scheme does not suffer from this difficulty.

Finally, technical difficulties with the present scheme should be pointed out. Since the present scheme utilizes single-particle entanglement and its multi-particle generalizations, it requires production, maintenance, and detection of photons at a single photon level. In particular, the Bell measurements required in the present scheme consist of identifying particular combinations of coincidence detection of a single photon among $(N+1)$ detectors. Although simple in principle, such Bell measurements require the highest degree of sensitivity on detectors, because detectors should be capable of distinguishing between no photon, one photon and two photons.

This research was supported by the Brain Korea 21 Project of the Korean Ministry of Education and by Korea Research Foundation. The author wishes to thank Professor J. Kim for stimulating discussions and Dr. D. H. Kwon




Figure 1

FIG. 1: Teleportation of a single-particle entangled state. An unknown entangled state in the beams a_1 and a_2 is teleported to the state in the beams B_1 and B_2 . The wavy line represents a classical communication channel.

for technical assistance in figures.

-
- [1] See, for example, The Physics of Quantum Information, edited by D. Bouwmeester, A. Ekert, and A. Zeilinger (Springer-Verlag, Berlin, 2000).
 - [2] C. H. Bennett, G. Brassard, C. Crepeau, R. Jozsa, A. Peres, and W. K. Wootters, Phys. Rev. Lett. **70**, 1895 (1993).
 - [3] D. Bouwmeester, J. W. Pan, K. Mattle, M. Eibl, H. Weinfurter, and A. Zeilinger, Nature (London) **390**, 575 (1997); D. Boschi, S. Branca, F. De Martini, L. Hardy, and S. Popescu, Phys. Rev. Lett. **80**, 1121 (1998); A. Furusawa, J. L. Sorensen, S. L. Braunstein, C. A. Fuchs, H. J. Kimble, and E. S. Polzik, Science **282**, 706 (1998).
 - [4] A. K. Ekert, Phys. Rev. Lett. **67**, 661 (1991).
 - [5] T. Jennewein, C. Simon, G. Weihs, H. Weinfurter, and A. Zeilinger, Phys. Rev. Lett. **84**, 4729 (2000); D. S. Naik, C. G. Peterson, A. G. White, A. J. Berglund, and P. G. Kwiat, *ibid* **84**, 4733 (2000); W. Tittel, J. Brendel, H. Zbinden, and N. Gisin, *ibid* **84**, 4737 (2000).
 - [6] A. Einstein, B. Podolsky, and N. Rosen, Phys. Rev. **47**, 777 (1935).
 - [7] J. S. Bell, Physics (Long Island City, N.Y.) **1**, 195 (1964).
 - [8] M. Zukowski, A. Zeilinger, M. A. Horne, and A. K. Ekert, Phys. Rev. Lett. **71**, 4287 (1993).
 - [9] S. Bose, V. Vedral, and P. L. Knight, Phys. Rev. A **57**, 822 (1998).
 - [10] J. W. Pan, D. Bouwmeester, H. Weinfurter, and A. Zeilinger, Phys. Rev. Lett. **80**, 3891 (1998).
 - [11] Here and throughout this paper, we assume that beam splitters are equipped with appropriate phase shifters such that two output modes are a symmetric and antisymmetric combination of two input modes.
 - [12] B. S. Shi, Y. K. Jiang, and G. C. Guo, Phys. Lett. A **268**, 161 (2000).
 - [13] L. Marinatto and T. Weber, Found. Phys. Lett. **13**, 119 (2000).
 - [14] D. M. Greenberger, M. A. Horne, A. Shimony, and A. Zeilinger, Amer. J. Phys. **58**, 1131 (1990); N. D. Mermin, Physics Today **43**, No. 6, 9 (1990).
 - [15] A. Zeilinger, M. A. Horne, H. Weinfurter, and M. Zukowski, Phys. Rev. Lett. **78**, 3031 (1997).
 - [16] H. W. Lee and J. Kim, Phys. Rev. A, in press.
 - [17] J. Lee and M. S. Kim, Phys. Rev. Lett. **84**, 4236 (2000).




Figure 2

FIG. 2: Setup at the source station for an "event-ready" experiment.




Figure 3

FIG. 3: Teleportation of a two-particle entangled state. An unknown entangled state in the beams a_1 , a_2 , b_1 , and b_2 is teleported to the state in the beams B_1 , B_2 , C_1 , and C_2 . The wavy line represents a classical communication channel.